

# INFLUENCE OF A SMALL PERTURBATION ON POINCARÉ-ANDRONOV OPERATORS WITH NOT WELL DEFINED TOPOLOGICAL DEGREE

OLEG MAKARENKOV<sup>1</sup>

(Submitted by J. Mawhin)

---

**ABSTRACT.** Let  $\mathcal{P}_\varepsilon \in C^0(\mathbf{R}^n, \mathbf{R}^n)$  be the Poincaré-Andronov operator over period  $T > 0$  of  $T$ -periodically perturbed autonomous system  $\dot{x} = f(x) + \varepsilon g(t, x, \varepsilon)$ , where  $\varepsilon > 0$  is small. Assuming that for  $\varepsilon = 0$  this system has a  $T$ -periodic limit cycle  $x_0$  we evaluate the topological degree  $d(I - \mathcal{P}_\varepsilon, U)$  of  $I - \mathcal{P}_\varepsilon$  on an open bounded set  $U$  whose boundary  $\partial U$  contains  $x_0([0, T])$  and  $\mathcal{P}_0(v) \neq v$  for any  $v \in \partial U \setminus x_0([0, T])$ . We give an explicit formula connecting  $d(I - \mathcal{P}_\varepsilon, U)$  with the topological indices of zeros of the associated Malkin's bifurcation function. The goal of the paper is to prove the Mawhin's conjecture claiming that  $d(I - \mathcal{P}_\varepsilon, U)$  can be any integer in spite of the fact that the measure of the set of fixed points of  $\mathcal{P}_0$  on  $\partial U$  is zero.

## 1. Introduction

Consider the system of ordinary differential equations

$$(1.1) \quad \dot{x} = f(x) + \varepsilon g(t, x, \varepsilon),$$

---

1991 *Mathematics Subject Classification.* Primary 54C40, 14E20; Secondary 46E25, 20C20.

*Key words and phrases.* Topological degree, perturbed Poincaré-Andronov map, zero measure singularities.

The work is partially supported by the Grant BF6M10 of Russian Federation Ministry of Education and CRDF (US), and by RFBR Grant 06-01-72552, and by the President of Russian Federation Young Candidate of Science grant MK-1620.2008.1.

where  $f \in C^1(\mathbf{R}^n, \mathbf{R}^n)$ ,  $g \in C^0(\mathbf{R} \times \mathbf{R}^n \times [0, 1], \mathbf{R}^n)$ ,  $g(t + T, v, \varepsilon) \equiv g(t, v, \varepsilon)$  and  $\varepsilon > 0$  is a small parameter. We suppose that equation (1.1) defines a flow in  $\mathbf{R}^n$ , i.e. assume the uniqueness and global existence for the solutions of the Cauchy problems associated to (1.1). For each  $v \in \mathbf{R}^n$  we denote by  $x_\varepsilon(\cdot, v)$  the solution of (1.1) with  $x_\varepsilon(0, v) = v$ . Thus, the Poincaré-Andronov operator over the period  $T > 0$  is defined by

$$\mathcal{P}_\varepsilon(v) := x_\varepsilon(T, v).$$

The problem of the existence (and even stability, see Ortega [11]) of  $T$ -periodic solutions of (1.1) with initial conditions inside an open bounded set  $U$  can be solved by evaluating the topological degree  $d(I - \mathcal{P}_\varepsilon, U)$  of  $I - \mathcal{P}_\varepsilon$  on  $U$  (see [6]). In the case when  $\mathcal{P}_0$  has no fixed points on the boundary  $\partial U$  of  $U$  the problem is completely solved by Capietto, Mawhin and Zanolin [2] who proved that  $d(I - \mathcal{P}_0, U) = (-1)^n d(f, U)$  generalizing the result by Berstein and Halanay [1] where  $U$  is assumed to be a neighborhood of an isolated zero of  $f$ . In the case when  $\mathcal{P}_0$  has fixed points on  $\partial U$  the pioneer result has been obtained by Mawhin [10] who considered the situation when  $f = 0$ . Mawhin proved that if  $g_0(v) = \int_0^T g(\tau, v, 0) d\tau$  does not vanish on  $\partial U$  then  $d(I - \mathcal{P}_\varepsilon, U)$  is defined for  $\varepsilon > 0$  sufficiently small and it can be evaluated as  $d(I - \mathcal{P}_\varepsilon, U) = d(-g_0, U)$ . This paper studies an intermediate situation when the fixed points of  $\mathcal{P}_0$  fill a part of  $\partial U$ . Current results on this subject deal with the case when  $\partial U$  contains a fixed number of fixed points, e.g. Feckan [4], Kamenskii-Makarenkov-Nistri [5]. As a part of a wider study of this problem Jean Mawhin (his seminar, November 2005) asked a question on evaluating  $d(I - \mathcal{P}_\varepsilon, U)$  in the case when  $\partial U$  contains a curve of fixed points of  $\mathcal{P}_0$ . He settled the following conjecture:

**Mawhin's conjecture.** *For small  $\varepsilon > 0$  the topological degree  $d(I - \mathcal{P}_\varepsilon, U)$  can be any integer depending on the perturbation term  $g$  in spite of the fact that the measure of  $\{v \in \partial U : \mathcal{P}_0(v) = v\}$  is zero.*

The goal of this paper is to evaluate  $d(I - \mathcal{P}_\varepsilon, U)$  and to give a proof of the above conjecture in the case when  $\{v \in \partial U : \mathcal{P}_0(v) = v\}$  forms a curve coming from a  $T$ -periodic limit cycle of the unperturbed system

$$(1.2) \quad \dot{x} = f(x).$$

Our fundamental assumption is that the algebraic multiplicity of the multiplier  $+1$  of the linearized system

$$(1.3) \quad \dot{y} = f'(x_0(t))y$$

equals to 1. In this case we say that the cycle  $x_0$  is nondegenerate.

The paper is organized as follows. In Section 2 for a fixed point  $v_\varepsilon$  of  $\mathcal{P}_\varepsilon$  satisfying  $v_\varepsilon \rightarrow v_0 \in x_0([0, T])$  as  $\varepsilon \rightarrow 0$  we obtain an asymptotic direction of the vector  $v_\varepsilon - v_0$ . By means of this result we evaluate in Section 3 the topological index of such fixed points  $v_\varepsilon \rightarrow v_0 \in x_0([0, T])$  as  $\varepsilon \rightarrow 0$  that  $v_\varepsilon \in U$ . Finally in Section 4 we give a proof of the Mawhin's conjecture provided that a technical assumption (see assumption 4.1) is satisfied.

## 2. Direction the fixed points of Poincaré-Andronov operator move when the perturbation increases

Since the cycle  $x_0$  is nondegenerate we can define (see [3], Ch. IV, § 20, Lemma 1) a matrix function  $Z_{n-1}$  solving the adjoint system

$$(2.1) \quad \dot{z} = -(f'(x_0(t)))^* z$$

and having the form  $Z_{n-1}(t) = \Phi(t)e^{\Lambda t}$ , where  $\Phi$  is a continuous  $T$ -periodic  $n \times n - 1$  matrix function and  $\Lambda$  is a  $n - 1 \times n - 1$ -matrix with different from 0 eigenvalues. Let  $z_0$  be the  $T$ -periodic solution of (2.1) satisfying  $z_0(0)^* \dot{x}_0(0) = 1$ . Finally, we denote by  $Y_{n-1}$  the  $n \times n - 1$  matrix function whose columns are solutions of the linearized system (1.3) satisfying  $Y_{n-1}(0)^* Z_{n-1}(0) = I$ .

The results of this paper are formulated in terms of the following auxiliary functions:

$$M(\theta) = \int_0^T z_0(\tau)^* g(\tau - \theta, x_0(\tau), 0) d\tau,$$

$$M^\perp(t, \theta) = (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t-T+\theta}^{t+\theta} (Z_{n-1}(\tau))^* g(\tau - \theta, x_0(\tau), 0) d\tau,$$

$$\angle(u, v) = \arccos \frac{\langle u, v \rangle}{\|u\| \cdot \|v\|}.$$

The function  $M$  was proposed by Malkin (see [9], formula 3.13) and the function  $M^\perp$  is a generalization of the function  $M_z^\perp$  of [8].

Next Theorem 2.1 shows that if a family  $\{x_{\varepsilon, \lambda}\}_{\lambda \in \Lambda}$  of  $T$ -periodic solutions of (1.1) emanate from  $x_0(\cdot + \theta_0)$  then a suitable projection of  $x_{\varepsilon, \lambda}(t) - x_0(t + \theta_0)$  can be always controlled. Though motivated by the Mawhin's conjecture, Theorem 2.1 can be of a general interest in the theory of oscillations playing a role of the first approximation formula (see Loud [7], formula 1.3, Lemma 1 and formula for  $x$  at p. 510) in the case when the zeros of the bifurcation function  $M$  are not necessary isolated.

**THEOREM 2.1.** *Let  $x_0$  be a nondegenerate  $T$ -periodic cycle of (1.2). Let  $\{x_{\varepsilon,\lambda}\}_{\lambda \in \Lambda}$  be a family of  $T$ -periodic solutions of (1.1) such that  $x_{\varepsilon,\lambda}(t) \rightarrow x_0(t + \theta_0)$  as  $\varepsilon \rightarrow 0$  uniformly with respect to  $t \in [0, T]$  and  $\lambda \in \Lambda$ . Then*

$$\angle(Z_{n-1}(t + \theta_0)^*(x_{\varepsilon,\lambda}(t) - x_0(t + \theta_0)), M^\perp(t, \theta_0)) \rightarrow 0 \quad \text{as } \varepsilon \rightarrow 0$$

*uniformly with respect to  $t \in [0, T]$  and  $\lambda \in \Lambda$ .*

**Proof.** The proof makes use of the idea of Theorem 3.1 of [8]. In the sequel  $(A, B)$  denotes the matrix composed by columns of matrixes  $A$  and  $B$ . Let  $a_\varepsilon \in C^0([0, T], \mathbf{R}^n)$  be given by

$$(2.2) \quad a_\varepsilon(t) = (z_0(t + \theta_0), Z_{n-1}(t + \theta_0))^*(x_\varepsilon(t) - x_0(t + \theta_0)).$$

Denoting  $Y(t) = (\dot{x}_0(t), Y_{n-1}(t))$  by Perron's lemma [12] (see also Demidovich ([3], Sec. III, §12) we have

$$(z_0(t), Z_{n-1}(t))^* Y(t) = I, \quad \text{for any } t \in \mathbf{R}.$$

Thus

$$(2.3) \quad x_\varepsilon(t) - x_0(t + \theta_0) = Y(t + \theta_0)a_\varepsilon(t), \quad \text{for any } t \in \mathbf{R}.$$

By subtracting (1.2) where  $x$  is replaced by  $x_0(\cdot + \theta_0)$  from (1.1) where  $x$  is replaced by  $x_\varepsilon$  we obtain

$$(2.4) \quad \begin{aligned} \dot{x}_\varepsilon(t) - \dot{x}_0(t + \theta_0) &= f'(x_0(t + \theta_0))(x_\varepsilon(t) - x_0(t + \theta_0)) \\ &+ \varepsilon g(t, x_\varepsilon(t), \varepsilon) + o(t, x_\varepsilon(t) - x_0(t + \theta_0)), \end{aligned}$$

where  $o(t, v)/\|v\| \rightarrow 0$  as  $\mathbf{R}^n \ni v \rightarrow 0$  uniformly with respect to  $t \in [0, T]$ . By substituting (2.3) into (2.4) we have

$$\begin{aligned} &\dot{Y}(t + \theta_0)a_\varepsilon(t) + Y(t + \theta_0)\dot{a}_\varepsilon(t) \\ &= f'(x_0(t + \theta_0))Y(t + \theta_0)a_\varepsilon(t) + \varepsilon g(t, x_\varepsilon(t), \varepsilon) + o(t, x_\varepsilon(t) - x_0(t + \theta_0)). \end{aligned}$$

Since  $f'(x_0(t))Y(t) = \dot{Y}(t)$  the last relation can be rewritten as

$$(2.5) \quad Y(t + \theta_0)\dot{a}_\varepsilon(t) = \varepsilon g(t, x_\varepsilon(t), \varepsilon) + o(t, x_\varepsilon(t) - x_0(t + \theta_0)).$$

Applying  $Z_{n-1}(t + \theta_0)^*$  to both sides of (2.5) we have

$$(0, I)\dot{a}_\varepsilon(t) = \varepsilon Z_{n-1}(t + \theta_0)^* g(t, x_\varepsilon(t), \varepsilon) + Z_{n-1}(t + \theta_0)^* o(t, x_\varepsilon(t) - x_0(t + \theta_0)),$$

where  $0$  denotes the  $n - 1$  dimensional zero vector and  $I$  stays for the identical  $n - 1 \times n - 1$  matrix. So

$$(0, I)a_\varepsilon(t) = (0, I)a_\varepsilon(t_0) + \varepsilon \int_{t_0}^t Z_{n-1}(\tau + \theta_0)^* g(\tau, x_\varepsilon(\tau), \varepsilon) d\tau$$

$$(2.6) \quad + \int_{t_0}^t Z_{n-1}(\tau + \theta_0)^* o(\tau, x_\varepsilon(\tau) - x_0(\tau + \theta_0)) d\tau.$$

From the definition of  $Z_{n-1}$  we have that  $Z_{n-1}(t)^* = (e^{\Lambda T})^* Z_{n-1}(t - T)^*$  for any  $t \in \mathbf{R}$  and so  $(0, I)a_\varepsilon(t)$  satisfies

$$(2.7) \quad (0, I)a_\varepsilon(t_0) = (e^{\Lambda T})^* (0, I)a_\varepsilon(t_0 - T) \quad \text{for any } t_0 \in [0, T].$$

Solving (2.6)-(2.7) with respect to  $(0, I)a_{\varepsilon, n}(t_0)$  we obtain

$$\begin{aligned} (0, I)a_\varepsilon(t_0) &= \varepsilon (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t_0-T}^{t_0} Z_{n-1}(\tau + \theta_0)^* g(\tau, x_\varepsilon(\tau), \varepsilon) d\tau \\ &+ (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t_0-T}^{t_0} Z_{n-1}(\tau + \theta_0)^* o(\tau, x_\varepsilon(\tau) - x_0(\tau + \theta_0)) d\tau \end{aligned}$$

for any  $t_0 \in [0, T]$ . On the other hand from (2.2) we obtain

$$Z_{n-1}(t + \theta_0)^* (x_\varepsilon(t) - x_0(t + \theta_0)) = (0, I)a_\varepsilon(t)$$

and therefore

$$(2.8) \quad \begin{aligned} &Z_{n-1}(t + \theta_0)^* (x_\varepsilon(t) - x_0(t + \theta_0)) - q_\varepsilon(t) \\ &= \varepsilon (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t-T}^t Z_{n-1}(\tau + \theta_0)^* g(\tau, x_\varepsilon(\tau), \varepsilon) d\tau, \end{aligned}$$

where

$$q_\varepsilon = (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t-T}^t Z_{n-1}(\tau + \theta_0)^* o(\tau, x_\varepsilon(\tau) - x_0(\tau + \theta_0)) d\tau.$$

From (2.8) we obtain

$$\begin{aligned} &\angle (Z_{n-1}(t + \theta_0)^* (x_\varepsilon(t) - x_0(t + \theta_0)), M^\perp(t, \theta_0)) \\ &= \angle \left( Z_{n-1}(t + \theta_0)^* \frac{x_\varepsilon(t) - x_0(t + \theta_0)}{\|x_\varepsilon - x_0(\cdot + \theta_0)\|_{[0, T]}}, M^\perp(t, \theta_0) \right) \\ &- \angle \left( Z_{n-1}(t + \theta_0)^* \frac{x_\varepsilon(t) - x_0(t + \theta_0)}{\|x_\varepsilon - x_0(\cdot + \theta_0)\|_{[0, T]}} - \frac{q_\varepsilon(t)}{\|x_\varepsilon - x_0(\cdot + \theta_0)\|_{[0, T]}}, M^\perp(t, \theta_0) \right) \\ &+ \angle \left( (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{t-T}^t Z_{n-1}(\tau + \theta_0)^* g(\tau, x_\varepsilon(\tau), \varepsilon) d\tau, M^\perp(t, \theta_0) \right). \end{aligned}$$

But the difference of the first two terms in the right hand part of the last equality tends to zero as  $\varepsilon \rightarrow 0$  and thus the thesis follows.

□

Next theorem is a reformulation of Theorem 2.1 suitable for our further considerations.

**THEOREM 2.2.** *Let  $x_0$  be a nondegenerate  $T$ -periodic cycle of (1.2). Let  $\{x_{\varepsilon,\lambda}\}_{\lambda \in \Lambda}$  be a family of  $T$ -periodic solutions of (1.1) such that  $x_{\varepsilon,\lambda}(t) \rightarrow x_0(t + \theta_0)$  as  $\varepsilon \rightarrow 0$  uniformly with respect to  $t \in [0, T]$  and  $\lambda \in \Lambda$ . Let  $l \in \mathbf{R}^n$  be an arbitrary vector such that  $\langle l, \dot{x}_0(\theta_0) \rangle = 0$ . Assume that  $\langle l, Y_{n-1}(\theta_0)M^\perp(0, \theta_0) \rangle \neq 0$ . Then there exists  $\varepsilon_0 > 0$  such that*

$$\langle l, x_{\varepsilon,\lambda}(0) - x_0(\theta_0) \rangle > 0 \quad \text{or} \quad \langle l, x_{\varepsilon,\lambda}(0) - x_0(\theta_0) \rangle < 0$$

according as

$$\langle l, Y_{n-1}(\theta_0)M^\perp(0, \theta_0) \rangle > 0 \quad \text{or} \quad \langle l, Y_{n-1}(\theta_0)M^\perp(0, \theta_0) \rangle < 0$$

for any  $\lambda \in \Lambda$  and any  $\varepsilon \in (0, \varepsilon_0]$ .

**Proof.** By Perron's lemma [12] (see also Demidovich ([3], Sec. III, §12) we have

$$v = Y_{n-1}(t)Z_{n-1}(t)^*v + \dot{x}_0(t)z_0(t)^*v$$

for any  $v \in \mathbf{R}^n$  and  $t \in \mathbf{R}$ . Therefore

$$\begin{aligned} & \langle l, x_{\varepsilon,\lambda}(0) - x_0(\theta_0) \rangle \\ &= \langle l, Y_{n-1}(\theta_0)Z_{n-1}(\theta_0)^*(x_{\varepsilon,\lambda}(0) - x_0(\theta_0)) \\ & \quad + \dot{x}_0(\theta_0)z_0(\theta_0)^*(x_{\varepsilon,\lambda}(0) - x_0(\theta_0)) \rangle \\ &= \langle Y_{n-1}(\theta_0)^*l, Z_{n-1}(\theta_0)^*(x_{\varepsilon,\lambda}(0) - x_0(\theta_0)) \rangle. \end{aligned}$$

Since  $\langle Y_{n-1}(\theta_0)^*l, M^\perp(0, \theta_0) \rangle \neq 0$  then by Theorem 2.1 there exists  $\varepsilon_0 > 0$  such that

$$\text{sign} \langle Y_{n-1}(\theta_0)^*l, Z_{n-1}(\theta_0)^*(x_{\varepsilon,\lambda}(0) - x_0(\theta_0)) \rangle = \text{sign} \langle Y_{n-1}(\theta_0)^*l, M^\perp(0, \theta_0) \rangle$$

for any  $\varepsilon \in (0, \varepsilon_0]$  and  $\lambda \in \Lambda$  and thus the proof is complete. □

### 3. The topological degree of the perturbed Poincaré-Andronov operator

To proceed to the proof of our main Theorem 3.1 we need three additional theorems which are formulated below for the convenience of the reader.

**Malkin's Theorem** (see [9], p. 41) *Assume that  $T$ -periodic solutions  $x_\varepsilon$  of (1.1) satisfy the property  $x_\varepsilon(t) \rightarrow x_0(t + \theta_0)$  as  $\varepsilon \rightarrow 0$ . Then  $M(\theta_0) = 0$ .*

**Capietto-Mawhin-Zanolin Theorem** (see [2], Corollary 2). *Let  $V \subset \mathbf{R}^n$  be an open bounded set. Assume that  $\mathcal{P}_0(v) \neq v$  for any  $v \in \partial V$ . Then  $d(I - \mathcal{P}_0, V) = (-1)^n d(f, V)$ .*

**Kamenskii-Makarenkov-Nistri Theorem** (see [5], Corollary 2.8). *Assume that  $\theta_0 \in [0, T]$  is an isolated zero of the bifurcation function  $M$ . Then there exist  $\varepsilon_0 > 0$  and  $r > 0$  such that  $\mathcal{P}_\varepsilon(v) \neq v$  for any  $\|v - v_0\| = r$  and any  $\varepsilon \in (0, \varepsilon_0]$ . Moreover  $d(I - \mathcal{P}_\varepsilon, B_r(v_0)) = \text{ind}(\theta_0, M)$ .*

We will say that the set  $U \subset \mathbf{R}^n$  has a smooth boundary if given any  $v \in \partial U$  there exists  $r > 0$  and a homeomorphism of  $\{\xi \in \mathbf{R}^{n-1} : \|\xi\| \leq 1\}$  onto  $\partial U \cap B_r(v)$ . Thus any set  $U$  with a smooth boundary possesses a tangent plane to  $\partial U$  at any  $v \in \partial U$ . This tangent plane will be denoted by  $L_U(v)$ . Moreover, if  $U$  has a smooth boundary and  $\mathbf{R}^n \ni h \notin L_U(v)$  then there exists  $\lambda_0 > 0$  such that either  $\lambda h + v \in U$  for any  $\lambda \in (0, \lambda_0]$  or  $\lambda h + v \notin U$  for any  $\lambda \in (0, \lambda_0]$ . In this case we will say that  $h$  centered at  $v$  is directed inward to  $U$  or outward respectively.

**THEOREM 3.1.** *Let  $x_0$  be a nondegenerate  $T$ -periodic cycle of (1.2). Let  $U \subset \mathbf{R}^n$  be an open bounded set with a smooth boundary and  $x_0([0, T]) \subset \partial U$ . Assume that  $\mathcal{P}_0(v) \neq v$  for any  $v \in \partial U \setminus x_0([0, T])$ . Assume that  $M$  has a finite number of zeros  $0 \leq \theta_1 < \theta_2 < \dots < \theta_k < T$  on  $[0, T]$  and  $\text{ind}(\theta_i, M) \neq 0$  for any  $i \in \overline{1, k}$ . Assume that  $Y_{n-1}(\theta_i)M^\perp(0, \theta_i) \notin L_U(x_0(\theta_i))$  for any  $i \in \overline{1, k}$ . Then for any  $\varepsilon > 0$  sufficiently small  $d(I - \mathcal{P}_\varepsilon, U)$  is defined. Moreover,*

$$d(I - \mathcal{P}_\varepsilon, U) = (-1)^n d(f, U) - \sum_{i=1}^k \text{ind}(\theta_i, M) D_i,$$

where  $D_i = 1$  or  $D_i = 0$  according as  $Y_{n-1}(\theta_i)M^\perp(0, \theta_i)$  centered at  $x_0(\theta_i)$  is directed inward to  $U$  or outward.

**Proof.** By Kamenskii-Makarenkov-Nistri theorem there exists  $r > 0$  and  $\varepsilon_0 > 0$  such that

$$(3.1) \quad d(I - \mathcal{P}_\varepsilon, B_r(x_0(\theta_i))) = \text{ind}(\theta_i, M)$$

for any  $\varepsilon \in (0, \varepsilon_0]$  and  $i \in \overline{1, k}$ . From Malkin's theorem we have the following "Malkin's property":  $r > 0$  can be decreased, if necessary, in such a way that there exists  $\varepsilon_0 > 0$  such that any  $T$ -periodic solution  $x_\varepsilon$  of (1.1) with initial condition  $x_\varepsilon(0) \in B_r(x_0([0, T]))$  and  $\varepsilon \in (0, \varepsilon_0]$  satisfies  $x_\varepsilon(0) \in \cup_{i \in \overline{1, k}} B_r(x_0(\theta_i))$ . Malkin's property implies that

$$(3.2) \quad d\left(I - \mathcal{P}_\varepsilon, \left(B_r(x_0([0, T])) \setminus \cup_{i \in \overline{1, k}} B_r(x_0(\theta_i))\right) \cap U\right) = 0$$

for any  $\varepsilon \in (0, \varepsilon_0]$ . Denote by  $l_i$  the perpendicular to  $L_U(x_0(\theta_i))$  directed outward away from  $U$  or inward according as  $(Z_{n-1}(\theta_i)^*)^{-1}M^\perp(0, \theta_i)$  centered at  $x_0(\theta_i)$  is directed outward away from  $U$  or inward. From Theorem 2.2 and Malkin's

property we have that  $\varepsilon_0 > 0$  can be diminished in such a way that for any  $i \in \overline{1, k}$  any  $T$ -periodic solution  $x_\varepsilon$  of (1.1) with initial condition  $x_\varepsilon(0) \in B_r(x_0(\theta_i))$  and  $\varepsilon \in (0, \varepsilon_0]$  satisfies  $x_\varepsilon(0) \in B_r(x_0(\theta_i)) \cap U$  or  $x_\varepsilon(0) \notin B_r(x_0(\theta_i)) \cap U$  according as  $D_i = 1$  or  $D_i = 0$ . This observation allows to deduce from (3.1) that

$$(3.3) \quad d(I - \mathcal{P}_\varepsilon, B_r(x_0(\theta_i)) \cap U) = \text{ind}(\theta_i, M), \quad \text{if } D(\theta_i) = 1,$$

$$(3.4) \quad d(I - \mathcal{P}_\varepsilon, B_r(x_0(\theta_i)) \cap U) = 0, \quad \text{if } D(\theta_i) = 0,$$

for any  $\varepsilon \in (0, \varepsilon_0]$  and  $i \in \overline{1, k}$ .

Observe that our choice of  $r > 0$  ensures that  $\mathcal{P}_0(v) \neq v$  for any  $v \in \partial(U \setminus B_r(x_0([0, T])))$ . Thus, by Capietto-Mawhin-Zanolin theorem we have  $d(I - \mathcal{P}_0, U \setminus B_r(x_0([0, T]))) = (-1)^n d(f, U \setminus B_r(x_0([0, T])))$ . Without loss of generality we can consider  $r > 0$  sufficiently small such that  $d(f, U \setminus B_r(x_0([0, T]))) = d(f, U)$  obtaining

$$(3.5) \quad d(I - \mathcal{P}_0, U \setminus B_r(x_0([0, T]))) = (-1)^n d(f, U).$$

Since

$$\begin{aligned} d(I - \mathcal{P}_\varepsilon, U) &= d\left(I - \mathcal{P}_\varepsilon, \left(B_r(x_0([0, T])) \setminus \bigcup_{i \in \overline{1, k}} B_r(x_0(\theta_i))\right) \cap U\right) \\ &\quad + d\left(I - \mathcal{P}_\varepsilon, \bigcup_{i \in \overline{1, k}} B_r(x_0(\theta_i)) \cap U\right) \\ &\quad + d(I - \mathcal{P}_\varepsilon, U \setminus B_r(x_0([0, T]))) \end{aligned}$$

the conclusion follows from formulas (3.2)-(3.5).  $\square$

#### 4. A proof of the Mawhin's conjecture

In this section we assume that the set  $U \subset \mathbf{R}^n$  has a smooth boundary and there exists  $v_{n-1} \in \mathbf{R}^{n-1}$  satisfying the following assumption

$$(4.1) \quad Y_{n-1}(t) (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} (e^{\Lambda t})^* v_{n-1} \notin L_U(t) \quad \text{for any } t \in [0, T].$$

We note that assumption (4.1) does not depend on the perturbation term of (1.1) and relies to unperturbed system (1.2). Let  $D = 1$  or  $D = 0$  according as  $Y_{n-1}(0) (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} (e^{\Lambda t})^* v_{n-1}$  centered at  $x_0(0)$  is directed inward to  $U$  or outward. Given odd  $m \in \mathbf{N}$  we construct the perturbation term  $g$  in such a way that  $d(I - \mathcal{P}_\varepsilon, U) = (-1)^n d(f, U) - m(2D - 1)$  for any  $\varepsilon > 0$  sufficiently small. Without loss of generality we consider  $T = 2\pi$ .

Since  $(z_0(t), Z_{n-1}(t))$  is nonsingular then  $((z_0(t), \Phi(t))^*)$  is nonsingular as well. Define  $\Omega : x_0([0, 2\pi]) \rightarrow \mathbf{R}^n$  as  $\Omega(x_0(t)) = ((z_0(t), \Phi(t))^*)^{-1}$  for any  $t \in [0, 2\pi]$ . By Uryson's theorem (see [6], Ch. 1, Theorem 1.1)  $\Omega$  can be continued to the whole  $\mathbf{R}^n$  in such a way that  $\Omega \in C^0(\mathbf{R}^n, \mathbf{R}^n)$ . Analogously, we consider



$\tilde{\Gamma} \in C^0(\mathbf{R}^n, \mathbf{R}^n)$  such that  $\tilde{\Gamma}(x_0(t)) = (\arcsin(\sin t), 0, \dots, 0)^*$  and denote by  $\Gamma \in C^0(\mathbf{R}^n, \mathbf{R})$  the first component of  $\tilde{\Gamma}$ . Let us define a  $2\pi$ -periodic  $\alpha$ -approximation of  $\left((e^{\Lambda t})^*\right)^{-1}$  on  $[-2\pi, 0]$  by

$$e_\alpha(t) = ((e^{\Lambda t})^*)^{-1}, \quad \text{if } t \in [-2\pi, -\alpha],$$

$$e_\alpha(t) = \frac{t}{-\alpha} \left((e^{-\Lambda\alpha})^*\right)^{-1} + \left(1 - \frac{t}{-\alpha}\right) \left((e^{-2\pi\Lambda})^*\right)^{-1}, \quad \text{if } t \in [-\alpha, 0],$$

which is continued to  $(-\infty, \infty)$  by the  $2\pi$ -periodicity. We are now in a position to introduce the required perturbation term, namely we consider that the perturbed system (1.1) has the following form

$$(4.2) \quad \dot{x} = f(x) + \varepsilon \Gamma(x) \Omega(x) \begin{pmatrix} D \sin(mt) + (1-D) \cos(mt) \\ (D \cos(mt) + (1-D) \sin(mt)) e_\alpha(t) v_{n-1} \end{pmatrix},$$

where  $\alpha > 0$  is sufficiently small. Consequently we denote by  $\mathcal{P}_\varepsilon$  the Poincaré-Andronov operator of system (4.2) over the period  $2\pi$ .

**PROPOSITION 4.1.** *Let  $x_0([0, T]) \subset U \subset \mathbf{R}^n$  be an open bounded set with a smooth boundary and assume that there exists  $v_{n-1} \in \mathbf{R}^n$  such that (4.1) is satisfied. Then given any odd  $m > 0$  there exists  $\alpha_0 > 0$  such that for any fixed  $\alpha \in (0, \alpha_0]$  and  $\varepsilon > 0$  sufficiently small  $d(I - \mathcal{P}_\varepsilon, U)$  is defined and*

$$d(I - \mathcal{P}_\varepsilon, U) = \begin{cases} (-1)^n d(f, U) - m, & \text{if } D = 1, \\ (-1)^n d(f, U) + m, & \text{if } D = 0. \end{cases}$$

**Proof.** By the definition of  $\Omega$  and  $\Gamma$  we have

$$(4.3) \quad \begin{pmatrix} z_0(t)^* \\ Z_{n-1}(t)^* \end{pmatrix} \Omega(x_0(t)) = \begin{pmatrix} 1 & 0 \\ 0 & (e^{\Lambda t})^* \end{pmatrix},$$

$$\Gamma(x_0(t)) = \arcsin(\sin t).$$

Therefore, taking into account that  $m$  is odd, we obtain the following formula for the bifurcation function  $M$

$$M(\theta) = \int_0^{2\pi} \arcsin(\sin \tau) (D \sin(m(\tau - \theta)) + (1-D) \cos(m(\tau - \theta))) d\tau$$

$$= (-1)^{(m-1)/2} \frac{4D \cos(m\theta) + 4(1-D) \sin(m\theta)}{m^2}$$

whose zeros are  $\theta_j = \frac{1}{m} \left( \frac{D\pi}{2} + j\pi \right)$ ,  $j \in \overline{0, 2m-1}$ . Moreover,

$$(4.4) \quad \text{ind}(\theta_j, M) = \text{sign}(M'(\theta_j))$$

$$= (-1)^{(m-1)/2} \text{sign} \left( \frac{4m \left( -D \sin \left( D \frac{\pi}{2} + j\pi \right) + (1-D) \cos \left( D \frac{\pi}{2} + j\pi \right) \right)}{m^2} \right).$$

Let us denote by  $M_\alpha^\perp$  the function  $M^\perp$  corresponding to system (4.2). From (4.3) we have that

$$\begin{aligned} M_\alpha^\perp(0, \theta) &= (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} \int_{-2\pi}^0 (Z_{n-1}(s + \theta))^* g(s, x_0(s + \theta), 0) ds \\ &= (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} (e^{\Lambda \theta})^* \circ \\ &\quad \circ \int_{-2\pi}^0 (e^{\Lambda s})^* e_\alpha(s) v_{n-1} \arcsin(\sin(s + \theta)) (D \cos(ms) + (1 - D) \sin(ms)) ds. \end{aligned}$$

Since

$$\begin{aligned} &\int_{-2\pi}^0 \arcsin(\sin(s + \theta)) (D \cos(ms) + (1 - D) \sin(ms)) ds \\ &= -(-1)^{(m-1)/2} \cdot \frac{4(D \sin(m\theta) + (1 - D) \cos(m\theta))}{m^2} \end{aligned}$$

by taking into account that  $m$  is odd we have that  $M_\alpha^\perp(0, \theta) \rightarrow M_0^\perp(0, \theta)$  as  $\alpha \rightarrow 0$ , where

$$\begin{aligned} M_0^\perp(0, \theta) &= - (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} (e^{\Lambda \theta})^* v_{n-1} (-1)^{(m-1)/2} \\ &\quad \cdot \frac{4(D \sin(m\theta) + (1 - D) \cos(m\theta))}{m^2}. \end{aligned}$$

Put  $q(\theta) = -(-1)^{(m-1)/2} (D \sin(m\theta) + (1 - D) \cos(m\theta))$ . Then, taking any  $\theta \in [0, 2\pi]$  and using the definition of  $D$  we conclude that  $Y_{n-1}(\theta) M_0^\perp(0, \theta)$  centered at  $x_0(\theta)$  is directed inward to  $U$  or outward according as  $\text{sign}(q(\theta))(2D - 1) = 1$  or  $\text{sign}(q(\theta))(2D - 1) = -1$ . Therefore, there exists  $\alpha_0 > 0$  such that for any  $\alpha \in [0, \alpha_0]$  and any  $\theta \in [0, 2\pi]$  we have that  $Y_{n-1}(\theta) M_\alpha^\perp(0, \theta)$  centered at  $x_0(\theta)$  is directed inward to  $U$  or outward according as  $\text{sign}(q(\theta))(2D - 1) = 1$  or  $\text{sign}(q(\theta))(2D - 1) = -1$ . Thus denoting by  $\mathcal{P}_{\varepsilon, \alpha}$  the Poincaré-Andronov operator of system (4.2) from Theorem 3.1 we have that

$$(4.5) \quad d(I - \mathcal{P}_{\varepsilon, \alpha}, U) = (-1)^n d(f, U) - \sum_{j \in \overline{0, 2m-1} : \text{sign}(q(\theta_j))(2D-1)=1} \text{ind}(\theta_j, M)$$

for any  $\alpha \in (0, \alpha_0]$ . Consider the case when  $D = 1$ . Then the property  $\text{sign}(q(\theta_j))(2D - 1) = 1$  is equivalent to

$$(4.6) \quad (-1)^{(m-1)/2} \text{sign}(\sin(\pi/2 + j\pi)) = -1.$$

If  $j \in \overline{0, 2m-1}$  satisfies (4.6) then (4.4) implies  $\text{ind}(\theta_j, M) = 1$ . Since there exists exactly  $m$  elements of  $\overline{0, 2m-1}$  satisfying (4.6) then (4.5) can be rewritten as  $d(I - \mathcal{P}_\varepsilon, U) = d(f, U) - m$ . Analogously, if  $D = 0$  then  $\text{sign}(q(\theta_j))(2D - 1) = 1$

is equivalent to  $(-1)^{(m-1)/2} \text{sign}(\cos(j\pi)) = -1$  that in combination with (4.4) gives  $\text{ind}(\theta_j, M) = -1$  allowing to rewrite (4.5) in the form  $d(I - \mathcal{P}_\varepsilon, U) = d(f, U) + m$ .

□

At the end of the paper we note that system (1.2) should exhibit very complex behavior in order that assumption (4.1) be not satisfied with any  $v_{n-1} \in \mathbf{R}^{n-1}$ . Particularly, (4.1) holds true for the prototypic unperturbed system (1.2)

$$(4.8) \quad \begin{aligned} \dot{x}_1 &= x_2 - x_1(x_1^2 + x_2^2 - 1), \\ \dot{x}_2 &= -x_1 - x_2(x_1^2 - x_2^2 - 1), \end{aligned}$$

$$\dot{x}_3 = -x_3$$

possessing the nondegenerate  $2\pi$ -periodic cycle  $x_0(t) = \begin{pmatrix} \sin t \\ \cos t \end{pmatrix}$  and  $U = B_1(0) = \{v \in \mathbf{R}^3 : \|v\| < 1\}$ . Indeed, it can be easily checked that  $\Phi(t) = \left( \begin{pmatrix} \sin t \\ 0 \end{pmatrix}, \begin{pmatrix} \cos t \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 1 \end{pmatrix} \right)^*$ ,  $e^{\Lambda t} = \begin{pmatrix} e^{2t} & 0 \\ 0 & e^t \end{pmatrix}$  and  $Y_{n-1}(t) = \Phi(t)e^{-\Lambda t}$  in this case. Thus, taking  $v_{n-1} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$  we have

$$Y_{n-1}(t) (e^{\Lambda T})^* \left( (e^{\Lambda T})^* - I \right)^{-1} (e^{\Lambda t})^* v_{n-1} = \frac{e^{2t}}{e^{2t} - 1} (\sin t, \cos t, 0)^*.$$

This last vector centered at  $x_0(t)$  is perpendicular to  $\partial U$  for any  $t \in [0, 2\pi]$ .

## REFERENCES

- [1] I. BERSTEIN AND A. HALANAY, *The index of a critical point and the existence of periodic solutions to a system with small parameter*, Dokl. Akad. Nauk **111** (1956), 923–925.
- [2] A. CAPIETTO, J. MAWHIN AND F. ZANOLIN, *Continuation theorems for periodic perturbations of autonomous systems*, Trans. Amer. Math. Soc. **329** (1992), 41–72.
- [3] B. P. DEMIDOVICH, *Lectures on the mathematical theory of stability*, Izdat. Nauka, Moscow, 1967.
- [4] M. FECKAN, *Bifurcation of periodic solutions in differential inclusions*, Appl. Math. **42** (1997), 369–393.
- [5] M. KAMENSKII, O. MAKARENKO AND P. NISTRI, *A continuation principle for a class of periodically perturbed autonomous systems*, Math. Nachr. **281** (2008), 42–61.
- [6] M. A. KRASNOSEL'SKII AND P. P. ZABREYKO, *Geometrical methods of nonlinear analysis. Fundamental Principles of Mathematical Sciences*, vol. 263, Springer-Verlag, Berlin, 1984.
- [7] W. S. LOUD, *Periodic solutions of a perturbed autonomous system*, Ann. of Math. **70** (1959), 490–529.
- [8] O. MAKARENKO AND P. NISTRI, *On the rate of convergence of periodic solutions in perturbed autonomous systems as the perturbation vanishes*, Comm. Pure Appl. Anal. **7** (2008), 49–61.
- [9] I. G. MALKIN, *On Poincaré's theory of periodic solutions*, Akad. Nauk SSSR. Prikl. Mat. Meh. **13** (1949), 633–646.
- [10] J. MAWHIN, *Le Problème des Solutions Périodiques en Mécanique non Linéaire*, Thèse de doctorat en sciences, Université de Liège, published in Degré topologique et solutions

- périodiques des systèmes différentiels non linéaires, Bull. Soc. Roy. Sci. Liège, Vol. 38, 1969, pp. 308–398.
- [11] R. ORTEGA, *A criterion for asymptotic stability based on topological degree*, Proceedings of the First World Congress of Nonlinear Analysts, Tampa, 1992, pp. 383–394.
- [12] O. PERRON, *Die Ordnungszahlen der Differentialgleichungssysteme*, Math. Zeitschr **31** (1930), 748–766.

*Manuscript received October 10, 2007*

OLEG MAKARENKOV

Research Institute of Mathematics, Voronezh State University, 394006, Voronezh, Universitetskaja pl.1

*Current address:* Research Institute of Mathematics, Voronezh State University, 394006, Voronezh, Universitetskaja pl.1

*E-mail address:* omakarenkov@math.vsu.ru